

# Investigation of the Common Mycorrhizal Network Concept: Plant Growth Responses in Simulated Intercropping of a Legume and Grass under Water Stress

BEDE S. MICKAN<sup>1,2,3</sup>, MIRANDA HART<sup>4</sup>, ZAKARIA M. SOLAIMAN<sup>1,2</sup>, KADAMBOT H. M. SIDDIQUE<sup>2</sup>, SASHA N. JENKINS<sup>1,2</sup>, LYNETTE K. ABBOTT<sup>1,2</sup>

<sup>1</sup>UWA School of Agriculture and Environment (M079), The University of Western Australia, Perth WA 6009, Australia

<sup>2</sup>The UWA Institute of Agriculture (M082), The University of Western Australia, Perth WA 6009, Australia

<sup>3</sup>Richgro Garden Products, 203 Acourt Rd, Jandakot Western Australia 6164, Australia.

<sup>4</sup>Department of Biology, University of British Columbia Okanagan, 3187 University Way, Kelowna, BC V1V 1V7, Canada

It is widely claimed that the common mycorrhizal network (CMN) plays a significant role in facilitated transfer of nutrients between plants. This experiment investigated the role of a common mycorrhizal network between a C3 legume and a C4 grass under nutrient and water-limited conditions. Shoot mass of *Trifolium subterraneum* increased by almost 150% when grown in close proximity to *Panicum clandestinum* when the only possible connection between roots was via a common mycorrhizal network. Inter-species competition between *T. subterraneum* and *P. clandestinum* in low nutrient soil was observed. The soil bacterial community was similar for both *T. subterraneum* and *P. clandestinum*. Water-stress increased the relative abundance of Firmicutes and Actinobacteria and decreased the relative abundance of Proteobacteria especially when they were most likely to share a mycorrhizal connection. Water-stress decreased the putative abundance of N-cycling genes under *P. clandestinum*, but not under *T. subterraneum*. The competitiveness of *T. subterraneum* when grown adjacent to *P. clandestinum* corresponded with enhanced P acquisition and depletion of soil P under *P. clandestinum*. It was concluded that the most likely explanation for competitiveness of *T. subterraneum* was enhanced P acquisition via mycorrhizal hyphae from soil in the *P. clandestinum* soil chamber rather than direct P transfer from *P. clandestinum* via a common mycorrhizal network